

Preventive Care Plan

Are Preventive Care Services covered only when performed in-network?

Yes, these preventive services are only covered under the preventive care plan when performed by an in-network provider. Your plan includes access to one of the largest preferred provider organization (PPO) networks. Details for locating an in-network provider can be found in the PPO Provider Network section of this guide.

Covered Preventive Services for Adults

Screenings for:

- Abdominal aortic aneurysm (one-time screening for men of specified ages who have ever smoked)
- Alcohol misuse
- Blood pressure
- Cholesterol (for adults of certain ages or at higher risk)
- Colorectal cancer (for adults over 45)
- Depression
- Type 2 diabetes (for adults with high blood pressure)
- Hepatitis B (for virus infection in persons with high risk)
- Hepatitis C (for infection in persons at high risk) (one-time screening for HCV to adults born between 1945-1965)
- HIV (for all adults at higher risk)
- Lung Cancer (for adults age 55-80 with a 30-pack per year smoking history and who currently smoke or quit within the past 15 years)
- Obesity
- Tobacco use
- Syphilis (for all adults at higher risk)

Counseling for:

- Alcohol misuse
- Aspirin use for men and women of certain ages and cardiovascular risk factors
- Diet (for adults with higher risk for chronic disease)
- Human Immunodeficiency Virus (HIV) for sexually active women
- Obesity
- Sexually transmitted infection (STI) prevention (for adults at higher risk)
- Tobacco use (including programs to help you stop using tobacco)

Immunizations:

- Doses, recommended ages, and recommended populations vary.
- Diphtheria, pertussis, tetanus (DPT)
- Hepatitis A
- Hepatitis B
- Herpes zoster
- Human papillomavirus (HPV)
- Influenza (Flu)
- Measles, mumps, rubella (MMR)
- Meningococcal (meningitis)
- Pneumococcal (pneumonia)
- Varicella (chicken pox)

Additional Covered Preventive Services for Women

- Aspirin (low dose as preventive after 12 weeks gestation in women who are at high risk for preeclampsia)
- Breast Cancer preventive medications for women with increased risk (tamoxifen or raloxifene).
- Contraception (FDA approved and ACA required contraceptive methods, sterilization procedures, and patient education and counseling)
- Well-woman visits (to obtain recommended preventive services for women under 65)

Screenings for:

- Breast cancer (mammography every 1 to 2 years for women over 40)
- Cervical cancer (for sexually active women)
- Chlamydia infection (for younger women and other women at higher risk)
- Domestic and interpersonal violence
- Gestational diabetes (for those at high risk)
- Gonorrhea (for all women at higher risk)
- Human Immunodeficiency Virus (HIV) (for sexually active women)
- Human Papillomavirus (HPV) DNA Test: High risk HPV DNA testing every three years for women with normal cytology results who are 30 or older
- Syphilis (for all pregnant women or other women at increased risk)
- Osteoporosis (for women over age 60 depending on risk factors)